

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 14/435, C12N 1/00, 1/10, 5/10, 15/12, 15/63</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/34326</b> <b>(43) International Publication Date:</b> 15 June 2000 (15.06.00)
<b>(21) International Application Number:</b> PCT/US99/29473 <b>(22) International Filing Date:</b> 10 December 1999 (10.12.99)  <b>(30) Priority Data:</b> 09/210,330 11 December 1998 (11.12.98) US 09/418,529 9 December 1999 (09.12.99) US  <b>(71) Applicant (for all designated States except US):</b> CLONTECH LABORATORIES, INC. [US/US]; 1020 East Meadow Drive, Palo Alto, CA 94303 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LUKYANOY, Sergey Anatolievich [RU/RU]; ul. Golubinskaya 13/1-161, Moscow (RU). FRADKOV, Arcady Fedorovich [RU/RU]; ul. Dnepropetrovskaya, 35/2-14, Moscow, 113570 (RU). LABAS, Yulii Aleksandrovich [RU/RU]; ul. Generala Tyuleneva, 35-416, Moscow, 117465 (RU). MATZ, Mikhail Vladimirovich [RU/RU]; ul. Teplii stan 7/2-28, Moscow, 117465 (RU). GREEN, Gisele [US/US]; 4106 Thain Way, Sunnyvale, CA 94087 (US). CHEN, Ying [CN/US]; 680 Garland Avenue, #6, Sunnyvale, CA 94086 (US). DING, Li [CN/US]; 1352 Norman Drive, Sunnyvale, CA 94087 (US).		<b>(74) Agent:</b> ADLER, Benjamin, A.; McGregor & Adler, 8011 Candle Lane, Houston, TX 77071 (US).  <b>(81) Designated States:</b> JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND USES THEREOF  <b>(57) Abstract</b>  The present invention is directed to novel fluorescent proteins from non-bioluminescent organisms from the Class Anthozoa. Also disclosed are cDNAs encoding the fluorescent proteins.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5  
10

**FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES  
OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND  
USES THEREOF**

10

**BACKGROUND OF THE INVENTION**

Cross-reference to Related Application

15 This is a divisional application of U.S.S.N. 09/210,330 filed  
on December 11, 1998.

Field of the Invention

20 This invention relates to the field of molecular biology.  
More specifically, this invention relates to novel fluorescent proteins,  
cDNAs encoding the proteins and uses thereof.

Description of the Related Art

25 Fluorescence labeling is a particularly useful tool for  
marking a protein, cell, or organism of interest. Traditionally, a  
protein of interest is purified, then covalently conjugated to a  
fluorophore derivative. For *in vivo* studies, the protein-dye complex is  
then inserted into cells of interest using micropipetting or a method of  
reversible permeabilization. The dye attachment and insertion steps,  
however, make the process laborious and difficult to control. An

alternative method of labeling proteins of interest is to concatenate or fuse the gene expressing the protein of interest to a gene expressing a marker, then express the fusion product. Typical markers for this method of protein labeling include  $\beta$ -galactosidase, firefly luciferase and bacterial luciferase. These markers, however, require exogenous substrates or cofactors and are therefore of limited use for *in vivo* studies.

A marker that does not require an exogenous cofactor or substrate is the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria*, a protein with an excitation maximum at 395 nm, a second excitation peak at 475 nm and an emission maximum at 510 nm. GFP is a 238-amino acid protein, with amino acids 65-67 involved in the formation of the chromophore.

Uses of GFP for the study of gene expression and protein localization are discussed in detail by Chalfie et al. in *Science* 263 (1994), 802-805, and Heim et al. in *Proc. Nat. Acad. Sci.* 91 (1994), 12501-12504. Additionally, Rizzuto et al. in *Curr. Biology* 5 (1995), 635-642, discuss the use of wild-type GFP as a tool for visualizing subcellular organelles in cells, while Kaether and Gerdes in *Febs Letters* 369 (1995), 267-271, report the visualization of protein transport along the secretory pathway using wild-type GFP. The expression of GFP in plant cells is discussed by Hu and Cheng in *Febs Letters* 369 (1995), 331-334, while GFP expression in *Drosophila* embryos is described by Davis et al. in *Dev. Biology* 170 (1995), 726-729.

Crystallographic structures of wild-type GFP and the mutant GFP S65T reveal that the GFP tertiary structure resembles a barrel (Ormo et al., (1996) *Science* 273: 1392-1395; Yang, et al., (1996) *Nature Biotech* 14: 1246-1251). The barrel consists of beta sheets in a compact antiparallel structure, where, in the center, an alpha helix

containing the chromophore is shielded by the barrel. The compact structure makes GFP very stable under diverse and/or harsh conditions such as protease treatment, making GFP an extremely useful reporter in general. However, the stability of GFP makes it sub-optimal for  
5 determining short-term or repetitive events.

A great deal of research is being performed to improve the properties of GFP and to produce GFP reagents useful and optimized for a variety of research purposes. New versions of GFP have been developed, such as a "humanized" GFP DNA, the protein product of  
10 which has increased synthesis in mammalian cells (Haas, et al., *Current Biology* 6 (1996), 315-324; Yang, et al., *Nucleic Acids Research* 24 (1996), 4592-4593). One such humanized protein is "enhanced green fluorescent protein" (EGFP). Other mutations to GFP have resulted in blue-, cyan- and yellow-green light emitting versions. Despite the great  
15 utility of GFP, however, other fluorescent proteins with properties similar to or different from GFP would be useful in the art. Novel fluorescent proteins result in possible new colors, or produce pH-dependent fluorescence. Other benefits of novel fluorescent proteins include fluorescence resonance energy transfer (FRET) possibilities  
20 based on new spectra and better suitability for larger excitation.

The prior art is deficient in novel fluorescent proteins wherein the DNA coding sequences are known. The present invention fulfills this long-standing need in the art.

## SUMMARY OF THE INVENTION

The present invention is directed to DNA sequences encoding fluorescent proteins selected from the group consisting of:

(a) an isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) an isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in  
5 codon sequence due to the degeneracy of the genetic code and that encodes a fluorescent protein. Preferably, the DNA is isolated from a non-bioluminescent organism from Class Anthozoa. More preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57, 62, 64 and 66 and the fluorescent protein has the amino  
10 acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory  
15 elements necessary for expression of the DNA in the cell. Preferably, the DNA encodes a fluorescent protein having the amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

In still another embodiment of the present invention, there is provided a host cell transfected with a vector of the present  
20 invention, such that the host cell expresses a fluorescent protein. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells, insect cells and yeast cells. A representative example of bacterial cell is an *E. coli* cell.

The present invention is also directed to an isolated and  
25 purified fluorescent protein coded for by DNA selected from the group consisting of: (a) isolated DNA from an organism from Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in

codon sequence due to the degeneracy of the genetic code, and which encodes a fluorescent protein. Preferably, the protein has the amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

5           The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of: (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which  
10       hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Corallimorpharia. More preferably,  
15       the organism is from Family Discosomatidae, Genus Discosoma. Most particularly, the present invention is drawn to a novel fluorescent protein from *Discosoma* sp. "red", drFP583, or a fusion fluorescent protein drFP583/dmFP592.

          The present invention is further directed to an amino acid  
20       sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein the sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group  
25       consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.

          Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the modified strategy of 3'-RACE used to isolate the target fragments. Sequences of the oligonucleotides used are shown in Table 2. Dp1 and Dp2 are the degenerate primers used in the first and second PCR, respectively (see Tables 3 and 4 for the sequences of degenerate primers). In the case of *Discosoma sp. "red"*, the first degenerate primer used was NGH (SEQ ID No. 4), and the second degenerate primer used was GEGa (SEQ ID No. 6) or NFP (SEQ ID No. 13) or PVMb (SEQ ID No. 16).

Figure 2 shows the excitation and emission spectrum of the novel fluorescent protein from *Discosoma sp. "red"*, drFP583.

Figure 3 shows separate detection of drFP583 and enhanced green fluorescent protein (EGFP) transiently expressed in HeLa cells. Figure 3A shows the image obtained for EGFP using Chroma filter set 31001. Figure 3B shows the image taken from the same field of view for drFP583 using filter set 31002. The images shown in Figures 3A and 3B were pseudocolored and overlayed in Figure 3C. Figure 3D shows phase contrast taken from the same field of view as that in Figure 3C.

Figure 4 shows separate detection of humanized drFP583, enhanced cyan fluorescent protein (ECFP) and enhanced yellow-green fluorescent protein (EYFP) in a triple-labeled HeLa cell. Figure 4A shows the image obtained from one cell co-expressing all three fluorescent proteins using Omega filter set XF 35 for DsRed1-Mito (drFP583-Mito, humanized drFP583). Figure 4B shows the image for EYFP-Tub using filter set XF 104. Figure 4C shows the image for ECFP-Nuc using filter set XF 114. The images shown in Figures 4A, 4B and 4C were pseudocolored and overlayed to show all three signals in one image in Figure 4D.



## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "GFP" refers to the basic green fluorescent protein from *Aequorea victoria*, including prior art versions of GFP engineered to provide greater fluorescence or fluoresce in different colors. The sequence of *Aequorea victoria* GFP (SEQ ID No. 54) has been disclosed in Prasher et al., *Gene* 111 (1992), 229-33.

As used herein, the term "EGFP" refers to mutant variant of GFP having two amino acid substitutions: F64L and S65T (Heim et al., *Nature* 373 (1995), 663-664). The term "humanized" refers to changes made to the GFP nucleic acid sequence to optimize the codons for expression of the protein in human cells (Yang et al., *Nucleic Acids Research* 24 (1996), 4592-4593).

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds. (1984)); "Animal Cell Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not  
5 limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed  
10 under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA  
15 sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

As used herein, the term "hybridization" refers to the process of association of two nucleic acid strands to form an  
20 antiparallel duplex stabilized by means of hydrogen bonding between residues of the opposite nucleic acid strands.

The term "oligonucleotide" refers to a short (under 100 bases in length) nucleic acid molecule.

"DNA regulatory sequences", as used herein, are  
25 transcriptional and translational control sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may be used to drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell

line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example, heterologous DNA includes coding sequence in a construct where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a coding sequence attached to heterologous promoter or enhancer elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

The amino acids described herein are preferred to be in the "L" isomeric form. The amino acid sequences are given in one-letter code (A: alanine; C: cysteine; D: aspartic acid; E: glutamic acid; F: phenylalanine; G: glycine; H: histidine; I: isoleucine; K: lysine; L: leucine; M: methionine; N: asparagine; P: proline; Q: glutamine; R: arginine; S: serine; T: threonine; V: valine; W: tryptophane; Y: tyrosine; X: any residue). NH<sub>2</sub> refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243 (1969), 3552-59 is used.

The present invention is directed to an isolated DNA selected from the group consisting of: (a) isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to isolated DNA of (a) and  
5 which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which encodes a fluorescent protein. Preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57, 62, 64 and 66, and the fluorescent  
10 protein has the amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory  
15 elements necessary for expression of the DNA in the cell. Specifically, the DNA encodes a fluorescent protein having the amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

In still another embodiment of the present invention, there is provided a host cell transfected with the vector of the present  
20 invention, which expresses a fluorescent protein of the present invention. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells and insect cells. A representative example of bacterial cell is an *E. coli* cell.

The present invention is also directed to a DNA sequence  
25 encoding a fluorescent protein selected from the group consisting of: (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent

protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Corallimorpharia. More preferably, the organism is from Family Discosomatidae, Genus Discosoma. Most preferably, the organism is *Discosoma* sp. "red".

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) an isolated protein encoded by a DNA which encodes a fluorescent protein wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated protein encoded by a DNA which hybridizes to isolated DNA of (a); and (c) an isolated protein encoded by a DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code. Preferably, the isolated and purified fluorescent protein is drFP583, or a fusion fluorescent protein drFP583/dmFP592.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16 and is used as a primer in polymerase chain reaction. Alternatively, it can be used as a probe for hybridization screening of the cloned genomic or cDNA library.

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

5

### **EXAMPLE 1**

#### **Biological Material**

Novel fluorescent proteins were identified from several  
10 genera of Anthozoa which do not exhibit any bioluminescence but have fluorescent color as observed under usual white light or ultraviolet light. Six species were chosen (see Table 1).

**TABLE 1**Anthozoa Species Used in This Study

Species	Area of Origination	Fluorescent Color
Anemonia majano	Western Pacific	bright green tentacle tips
Clavularia sp.	Western Pacific	bright green tentacles and oral disk
Zoanthus sp.	Western Pacific	green-yellow tentacles and oral disk
Discosoma sp. "red"	Western Pacific	orange-red spots oral disk
Discosoma striata	Western Pacific	blue-green stripes on oral disk
Discosoma sp. "magenta"	Western Pacific	faintly purple oral disk
Discosoma sp. "green"	Western Pacific	green spots on oral disk
Anemonia sulcata	Mediterranean	purple tentacle tips

5

**EXAMPLE 2**cDNA Preparation

Total RNA was isolated from the species of interest  
 10 according to the protocol of Chomczynski and Sacchi (Chomczynski P.,



et al., *Anal. Biochem.* 162 (1987), 156-159). First-strand cDNA was synthesized starting with 1-3  $\mu$ g of total RNA using SMART PCR cDNA synthesis kit (CLONTECH) according to the provided protocol with the only alteration being that the "cDNA synthesis primer" provided in the  
5 kit was replaced by the primer TN3 (5'- CGCAGTCGACCG(T)<sub>13</sub>, SEQ ID No. 1) (Table 2). Amplified cDNA samples were then prepared as described in the protocol provided except the two primers used for PCR were the TS primer (5'-AAGCAGTGGTATCAACGCAGAGT, SEQ ID No. 2) (Table 2) and the TN3 primer (Table 2), both in 0.1  $\mu$ M concentration.  
10 Twenty to twenty-five PCR cycles were performed to amplify a cDNA sample. The amplified cDNA was diluted 20-fold in water and 1  $\mu$ l of this dilution was used in subsequent procedures.

**TABLE 2**Oligos Used in cDNA Synthesis and RACE

5 TN3: 5'-CGCAGTCGACCG(T)<sub>13</sub>  
(SEQ ID No. 1)

T7-TN3: 5'-GTAATACGACTCACTATAGGGCCGACGTCGACCG(T)<sub>13</sub>  
(SEQ ID No. 17)

10

TS-primer: 5'-AAGCAGTGGTATCAACGCAGAGT  
(SEQ ID No. 2)

15 T7-TS:  
5'-GTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT  
(SEQ ID No. 18)

T7: 5'-GTAATACGACTCACTATAGGGC  
(SEQ ID No. 19)

20

TS-oligo 5'-AAGCAGTGGTATCAACGCAGAGTACGCrGrGrG  
(SEQ ID No. 53)

**EXAMPLE 3**

25

Oligo Design

To isolate fragments of novel fluorescent protein cDNAs, PCR using degenerate primers was performed. Degenerate primers were designed to match the sequence of the mRNAs in regions that were predicted to be the most invariant in the family of fluorescent proteins. Four such stretches were chosen (Table 3) and variants of degenerate primers were designed. All such primers were directed to

30

the 3'-end of mRNA. All oligos were gel-purified before use. Table 2 shows the oligos used in cDNA synthesis and RACE.

**TABLE 3**

5

Key Amino Acid Stretches and Corresponding Degenerate Primers Used for Isolation of Fluorescent Proteins

Stretch Position according to A. victoria GFP (7)	Amino Acid Sequence of the Key Stretch	Degenerated Primer Name and Sequence
20-25	GXVNGH (SEQ ID No. 3)	NGH: 5'- GA(C,T) GGC TGC GT(A,T,G,C) AA(T,C) GG(A,T,G) CA (SEQ ID No. 4)
31-35	GEGEG (SEQ ID No. 5)  GEGNG (SEQ ID No. 8)	GEGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) GA(A,G) GG (SEQ ID No. 6) GEGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) GA(A,G) GG (SEQ ID No. 7) GNGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) AA(C,T) GG (SEQ ID No. 9) GNGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) AA(C,T) GG (SEQ ID No. 10)
127-131	GMNFP (SEQ ID No. 11) GVNFP (SEQ ID No. 12)	NFP: 5' TTC CA(C,T) GGT (G,A)TG AA(C,T) TT(C,T) CC (SEQ ID NO. 13)
134-137	GPVM (SEQ ID No. 14)	PVMa: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(A,C) ATG (SEQ ID NO. 15) PVMb: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(G,T) ATG (SEQ ID NO. 16)

#### EXAMPLE 4

##### Isolation of 3'-cDNA Fragments of nFPs

The modified strategy of 3'-RACE was used to isolate the target fragments (see Figure 1). The RACE strategy involved two consecutive PCR steps. The first PCR step involved a first degenerate primer (Table 4) and the T7-TN3 primer (SEQ ID No. 17) which has a 3' portion identical to the TN3 primer used for cDNA synthesis (for sequence of T7-TN3, Table 2). The reason for substituting the longer T7-TN3 primer in this PCR step was that background amplification which occurred when using the shorter TN3 primer was suppressed effectively, particularly when the T7-TN3 primer was used at a low concentration (0.1  $\mu$ M) (Frohman et al., (1998) *PNAS USA*, 85, 8998-9002). The second PCR step involved the TN3 primer (SEQ ID No. 1, Table 2) and a second degenerate primer (Table 4).

**TABLE 4**

Combinations of Degenerate Primers for First and Second PCR Resulting in Specific Amplification of 3'-Fragments of nFP cDNA

Species	First Degenerate Primer	Second Degenerate Primer
Anemonia majano	NGH (SEQ ID No. 4)	GNGb (SEQ ID No. 10)
Clavularia sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Zoanthus sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Discosoma sp. "red"	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6), NFP (SEQ ID No. 13) or PVMb (SEQ ID No. 16)
Discosoma striata	NGH (SEQ ID No. 4)	NFP (SEQ ID No. 13)
Anemonia sulcata	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6) or NFP (SEQ ID No. 13)

5

The first PCR reaction was performed as follows: 1 µl of 20-fold  
 10 dilution of the amplified cDNA sample was added into the reaction  
 mixture containing 1X Advantage KlenTaq Polymerase Mix with  
 provided buffer (CLONTECH), 200 µM dNTPs, 0.3 µM of first degenerate

primer (Table 4) and 0.1  $\mu$ M of T7-TN3 (SEQ ID No. 17) primer in a total volume of 20  $\mu$ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C, 1 min.; 72°C, 40 sec; 24 cycles for 95°C, 10 sec.; 62°C, 30 sec.; 72°C, 40 sec. The reaction was then diluted 20-fold in water and 1  $\mu$ l of this dilution was added to a second PCR reaction, which contained 1X Advantage KlenTaq Polymerase Mix with the buffer provided by the manufacturer (CLONTECH), 200  $\mu$ M dNTPs, 0.3  $\mu$ M of the second degenerate primer (Table 4) and 0.1  $\mu$ M of TN3 primer. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C (for GEG/GNG or PVM) or 52°C (for NFP), 1 min.; 72°C, 40 sec; 13 cycles for 95°C, 10sec.; 62°C (for GEG/GNG or PVM) or 58°C (for NFP), 30 sec.; 72°C, 40 sec. The product of PCR was cloned into PCR-Script vector (Stratagene) according to the manufacturer's protocol.

Different combinations of degenerate primers were tried in the first and second PCR reactions on the DNA from each species until a combination of primers was found that resulted in specific amplification--meaning that a pronounced band of expected size (about 650-800 bp for NGH and GEG/GNG and 350-500 bp for NFP and PVM--sometimes accompanied by a few minor bands) was detected on agarose gel after two PCR reactions. The primer combinations of choice for different species of the Class Anthozoa are listed in Table 4. Some other primer combinations also resulted in amplification of fragments of correct size, but the sequence of these fragments showed no homology to the other fluorescent proteins identified or to *Aequorea victoria* GFP.

## EXAMPLE 5

### Obtaining Full-Length cDNA Copies

Upon sequencing the obtained 3'-fragments of novel  
5 fluorescent protein cDNAs, two nested 5'-directed primers were  
synthesized for cDNA (Table 5), and the 5' ends of the cDNAs were  
then amplified using two consecutive PCRs. In the next PCR reaction,  
the novel approach of "step-out PCR" was used to suppress background  
amplification. The step-out reaction mixture contained 1x Advantage  
10 KlenTaq Polymerase Mix using buffer provided by the manufacturer  
(CLONTECH), 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the first gene-specific primer  
(see Table 5), 0.02  $\mu$ M of the T7-TS primer (SEQ ID No. 18), 0.1  $\mu$ M of  
T7 primer (SEQ ID No. 19) and 1  $\mu$ l of the 20-fold dilution of the  
amplified cDNA sample in a total volume of 20  $\mu$ l. The cycling profile  
15 was (Hybaid OmniGene Thermocycler, tube control mode): 23-27  
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of  
amplification was diluted 50-fold in water and one  $\mu$ l of this dilution  
was added to the second (nested) PCR. The reaction contained 1X  
Advantage KlenTaq Polymerase Mix with provided buffer (CLONTECH),  
20 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the second gene-specific primer and 0.1  $\mu$ M  
of TS primer (SEQ ID No. 2) in a total volume of 20  $\mu$ l. The cycling  
profile was (Hybaid OmniGene Thermocycler, tube control mode): 12  
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of  
amplification was then cloned into pAtlas vector (CLONTECH) according  
25 to the manufacturer's protocol.

**TABLE 5**Gene-Specific Primers Used for 5'-RACE

Species	First Primer	Second (Nested) Primer
Anemonia majano	5'-GAAATAGTCAGGCATACTGGT (SEQ ID No. 20)	5'-GTCAGGCATAC TGGTAGGAT (SEQ ID No. 21)
Clavularia sp.	5'-CTTGAAATAGTCTGCTATATC (SEQ ID No. 22)	5'-TCTGCTATATC GTCTGGGT (SEQ ID No. 23)
Zoanthus sp.	5'- GTTCTTGAAATAGTCTACTATGT (SEQ ID No. 24)	5'-GTCTACTATGTCTT GAGGAT (SEQ ID No. 25)
Discosoma sp. "red"	5'-CAAGCAAATGGCAAAGGTC (SEQ ID No. 26)	5'-CGGTATTGTGGCC TTCGTA (SEQ ID No. 27)
Discosoma striata	5'-TTGTCTTCTTCTGCACAAC (SEQ ID No. 28)	5'-CTGCACAACGG GTCCAT (SEQ ID No. 29)
Anemonia sulcata	5'-CCTCTATCTTCATTTCTGC (SEQ ID No. 30)	5'-TATCTTCATTTCTT GCGTAC (SEQ ID No. 31)
Discosoma sp. "magenta"	5'-TTCAGCACCCCATCACGAG (SEQ ID No. 32)	5'-ACGCTCAGAGCTG GGTTCC (SEQ ID No. 33)
Discosoma sp. "green"	5'-CCCTCAGCAATCCATCACGTTC (SEQ ID No. 34)	5'-ATTATCTCAGTGGA TGGTTC (SEQ ID No. 35)



## EXAMPLE 6

### Expression of nFPs in *E. coli*

To prepare a DNA construct for novel fluorescent protein expression, two primers were synthesized for each cDNA: a 5'-directed "downstream" primer with the annealing site located in the 3'-UTR of the cDNA and a 3'-directed "upstream" primer corresponding to the site of translation start site (not including the first ATG codon) (Table 6). Primers with SEQ ID Nos. 43 and 44 were the primers used to prepare the dr583 DNA. Both primers had 5'-heels coding for a site for a restriction endonuclease; in addition, the upstream primer was designed so as to allow the cloning of the PCR product into the pQE30 vector (Qiagen) in such a way that resulted in the fusion of reading frames of the vector-encoded 6xHis-tag and nFP. The PCR was performed as follows: 1 µl of the 20-fold dilution of the amplified cDNA sample was added to a mixture containing 1x Advantage KlenTaq Polymerase Mix with buffer provided by the manufacturer (CLONTECH), 200 µM dNTPs, 0.2 µM of upstream primer and 0.2 µM of downstream primer, in a final total volume of 20 µl. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 23-27 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of this amplification step was purified by phenol-chlorophorm extraction and ethanol precipitation and then cloned into pQE30 vector using restriction endonucleases corresponding to the primers' sequence according to standard protocols.

All plasmids were amplified in XL-1 blue *E. coli* and purified by plasmid DNA miniprep kits (CLONTECH). The recombinant clones were selected by colony color, and grown in 3 ml of LB medium (supplemented with 100 µg/ml of ampicillin) at 37°C overnight. 100 µl

of the overnight culture was transferred into 200 ml of fresh LB medium containing 100 µg/ml of ampicillin and grown at 37°C, 200 rpm up to OD<sub>600</sub> 0.6-0.7. 1 mM IPTG was then added to the culture and incubation was allowed to proceed at 37°C for another 16 hours. The  
5 cells were harvested and recombinant protein, which incorporated 6x His tags on the N-terminus, was purified using TALON™ metal-affinity resin according to the manufacturer's protocol (CLONTECH).

TABLE 6

Primers Used to Obtain Full Coding Region of nFPs for Cloning into Expression Construct

Species	Upstream Primer	Downstream Primer
Anemonia majano	5' -acatggatccgctctttcaaaca agttatc (SEQ ID No. 36) BamHI	5'-tagtactcgcgagcttattcgta tttcagtgaatc (SEQ ID No. 37) XhoI
Clavularia sp.	L: 5'-acatggatccaacattttttga gaaacg (SEQ ID No. 38) BamHI S: 5'-acatggatccaaagctctaacc accatg (SEQ ID No. 39) BamHI	5'-tagtactcgcgagcaacacaa accctcagacaa (SEQ ID No. 40) XhoI
Zoanthus sp.	5'- acatggatccgctcagtc aaag cacggt (SEQ ID No. 41) BamHI	5'-tagtactcgcgaggttggaactacat tcttatca (SEQ ID No. 42) XhoI
Discosoma sp. "red"	5'- acatggatccaggtcttccaagaat gttatc (SEQ ID No. 43) BamHI	5'-tagtactcgcgaggagccaagtc agcctta (SEQ ID No. 44) XhoI
Discosoma striata	5'- acatggatccagttgggtccaagagtgtg (SEQ ID No. 45) BamHI	5'-tagcgcgagctctatcatgcctc gtcacct (SEQ ID No. 46) SacI
Anemonia sulcata	5'- acatggatccgcttccttttaagaagact (SEQ ID No. 47) BamHI	5'-tagtactcgcgagtccttgggagc ggcttg (SEQ ID No. 48) XhoI
Discosoma sp. "magenta"	5'- acatggatccagttgttccaagaatgtgat (SEQ ID No. 49) BamHI	5'-tagtactcgcaggccattacg ctaatac (SEQ ID No. 50) XhoI
Discosoma sp. "green"	5'-acatggatccagtgcaactaaagaagaatg (SEQ ID No. 51)	5'-tagtactcgcgagattcggtttaat gccttg (SEQ ID No. 52)

**EXAMPLE 7****Novel Fluorescent Protein drFP583 and cDNA Encoding drFP583**

One of the full-length cDNAs encoding fluorescent proteins found is described herein (drFP583). The nucleic acid sequence and deduced amino acid sequence are SEQ ID Nos. 55 and 56, respectively. The spectral properties of drFP583 is listed in Table 7, and the emission and excitation spectra for the drFP583 is shown in Figure 2.

**TABLE 7****Spectral Properties of the Isolated drFP583**

Species:	Discosoma sp. "red"	Max. Extinction Coefficient:	22,500
nFP Name:	drFP583	Quantum Yield	0.23
Absorbance Max. (nm):	558	Relative Brightness:*	0.24
Emission Max. (nm):	583		

\*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

**EXAMPLE 8****Expression of drFP583 in Mammalian Cells**

HeLa cells were transfected either with plasmid pDsRed1-N1 (vector containing the DNA encoding drFP583) or plasmid pEGFP-C1

(encoding EGFP from *Aequoria victoria*). Immediately after the transfection, cells were mixed and plated on coverslips. Cells were incubated for 48 hours at 37°C then fixed in 3.7 % formaldehyde. Cells were mounted in mounting medium and observed by fluorescence  
5 microscopy. Images were taken from the same field of view with Chroma filter set 31001 for EGFP (Figure 3A), and filter set 31002 for drFP583 (Figure 3B) using a cooled CCD camera (Roper Scientific) and MetaMorph software (Universal Imaging). The images (Figures 3A and 3B) were pseudocolored and overlayed in Figure 3C. Phase contrast was  
10 taken from the same field of view and overlayed in Figure 3D.

## EXAMPLE 9

### Generation of Humanized drFP583

Since mammalian expression is a very popular tool, human  
5 favored codon version is needed for better expression in mammalian  
cells. Humanized drFP583 was therefore generated by changing wild  
type drFP583 nucleotide sequence to optimize the codons for  
expression of the fluorescent protein. The nucleotide sequence of the  
humanized drFP583 is shown in SEQ ID No. 57.

10

## EXAMPLE 10

### Expression of Humanized drFP583 in Mammalian Cells

15 HeLa cells were transiently co-transfected with plasmids  
pECFP-Nuc, pEYFP-Tub and pDsRed1-Mito (humanized drFP583). After  
transfection, cells were incubated for 48 hours at 37 °C then fixed in 3.7  
% formaldehyde. Cells were mounted in mounting medium and  
observed by fluorescence microscopy. Images were taken of one cell  
20 co-expressing all three fluorescent proteins with Omega filter set XF 35  
for DsRed1-Mito (Figure 4A), XF 104 for EYFP-Tub (Figure 4B) and XF  
114 for ECFP-Nuc (Figure 4C) using a cooled CCD camera (Roper  
Scientific) and MetaMorph software (Universal Imaging). Individual  
images were pseudocolored and overlaid to show all three signals in  
25 one image (Figure 4D). Protein DsRed1-Mito localizes to mitochondria,  
EYFP-Tub localizes to the microtubular network, and ECFP-Nuc localizes  
to the nucleus.

As a conclusion, drFP583 does emit to a low extent also in  
the cyan (ECFP), green (EGFP) and yellow-green (EYFP) emission channels

(filter sets). High expression levels or highly concentrated protein in intracellular structures can therefore result in high signal intensities that will give some bleedthrough in the other emission wavelengths. The bleedthrough is small and should not affect multiple labeling  
5 recording in most cases.

### **EXAMPLE 11**

#### 10 **Mutants of Humanized drFP583**

Mutants of humanized drFP583 were generated using error prone PCR technique (Clontech). Mutations occurred at amino acids 42, 71, 105, 120, 161 and 197 (numbering starting from the first Methionine). Table 8 lists the mutants that were generated and their  
15 properties.

TABLE 8

Mutants of Humanized drFP583

5

Mutant	Mutations	Properties
E5	V105A, S197T	Overnight in <i>E. coli</i> emitting green fluorescence; <i>in vitro</i> maturing to red over at 28h 37°C on 80% (retains 20% green peak); folding faster than wild type drFP583 (~28h)
E8	N42H	always two peaks green & red (~1:1) folding faster than E5 (~8h)
E5up	V105A	red from the beginning; folding faster than E5 (~12h)
E5down	S197T	phenotype is similar to E5
E57	V105A, I161, S197A	like E5 but folding faster (~8-10h); ~5% of green peak left at the end
AG4	V71M, V105A, S197T	bright green, no red at all; fast folding (~ 16h)
AG4H	V71M, V105A, Y120H, S197T	like AG4 but twice brighter



## EXAMPLE 12

### Characterization and Applications of E5 Mutant

5 E5 (V105A, S197T) changes its fluorescence from green to red over time both *in vitro* and *in vivo*, in *E. coli* and in mammalian cells. Also, E5 folds faster than wild type drFP583 both in *E. coli* and mammalian cells (Table 8).

Since it allows the "two color" reporting mode for  
10 monitoring of the promoter activity, i.e., for both active or shutdown state of the promoter, there is a separate color, serving as an indicator of that state, E5 can be used as a transcriptional reporter. Different from "one color" mode, "two color" mode has a measurable signal (color) present for both states of the process as opposed to "one  
15 color" mode (e.g. destabilized GFP) wherein the absence of the color is an indicator of the second state. Namely, newly produced E5 protein fluoresces in green, indicating on-going promoter activity. Over time, the protein will mature, acquiring the red fluorescence. So if the promoter is no longer active, all the protein will eventually mature,  
20 resulting in the dominant red fluorescence. In case the promoter is still active both red and green fluorescence will be readily detected. Thus E5 as a "two color" reporter allows study of gene expression similar to destabilized GFP, but with permanent "signature" of past gene activity in the cells, tissues or entire organism. For example, at the  
25 tissue level, E5 may help to distinguish the stem cells from differentiated cells. Providing the promoter is only active in the stem cells, the E5 reporter will label the stem cell population in green and red, the progenitor cells would be labeled predominantly in red, the terminally differentiated will not fluoresce (due to the titration out of  
30 protein during cell division).

E5 could be used for spatial and temporal visualization of newly synthesized vs. accumulated fusion proteins. That is, E5 could function like a fusion tag. Possible applications envisaged at different organizational levels. At the cellular level, E5 may help to visualize and  
5 distinguish the newly synthesized proteins in various compartments such as outer membrane, microvillae, ER, Golgi, mitochondria, nuclei, various components of cellular matrix and focal adhesion complexes. At the tissue level, E5 may be helpful in visualizing newly formed vs. preexisting structures e.g. membrane junctions, components of  
10 extracellular matrix.

One of the most fascinating applications of E5 seems to be in the study of the mother-daughter relationship during the cell division and migration. A wide horizon is opening in the fields of development as well as in the adult organisms to study the cell  
15 migration and differentiation. Allowing visualization of the expression "history", E5 can help to distinguishing between the mother cells where the promoter is actually active vs. the daughter cells containing the accumulated protein but not producing fresh protein anymore. It would enable the study of the cell fates during development and organ  
20 remodeling, thus distinguishing between cell migration and cell expansion or differentiation.

In conclusion, E5 is basically applicable to any situation where GFP was previously used. Main advantage is that E5 can track down "the history" of promoter activity or protein localization in cells  
25 or tissues. With a better protein stability than GFP, E5 allows longer analysis window (wild type drFP583 is stable for at least 4 weeks in *Xenopus*, while EGFP starts to faint after two weeks).

### **EXAMPLE 13**

#### **Characterization and Applications of E8 Mutant**

E8 (N42H) has two fluorescence maximums, green and red,  
5 at all times and it folds much faster than drFP583 (Table 8).

Since it detects both green and red fluorescence simultaneously, E8 may be useful for studying processes related to blood circulation and proteins/cells trafficking. Blood absorbs the green fluorescence; thus only the red fluorescence will be visible while  
10 the protein is trafficking in the blood. Both green and red fluorescence could be detected outside the bloodstream making the whole process easy to visualize and record. Monitoring both red and green fluorescence simultaneously may also help to reduce the background fluorescence problems for some tissues or cells.

15

### **EXAMPLE 14**

#### **Generation of drFP583/dmFP592 Hybrid Using Shuffling Procedure**

Non-humanized wild type coding region fragments from  
20 drFP583 and dmFP592 were amplified by PCR (22 cycles, 95°C, 15 sec., 68°C 1 min 20 sec.) using 1 ng of corresponding bacterial expression plasmids (pQE-30 derivatives with drFP583 and dmFP592 inserts, respectively) as templates. Oligonucleotides A (ACATGG  
25 ATCCAGGTCTTCCAAGAATGTTATC, SEQ ID No. 58), B (TAGTACTCG AGCCAAGTTCAGCCTTA, SEQ ID No. 59), C (ACATGGATCCAG TTGTTC CAAGAATGTGAT, SEQ ID No. 60), and D (TAGTACTCGAGGCCATTA CCGCTAATC, SEQ ID No. 61) were used as primers for amplifying these fragments in a concentration of 0.2 mM.

The PCR products were then purified by QIAquick PCR Purification Kit (QIAGEN). Afterwards, the purified fragments drFP583 and dmFP592 (300-500 ng each) were digested with restriction endonucleases EcoRI, HindIII and DraI (10 U each) simultaneously. Reactions were performed in BamHI restriction buffer (NEB) supplemented BSA for 3h at 37°C. Total reaction volume was 30 ml. Upon completion, the resulted restriction fragments from each restriction reaction were separated by electrophoresis in agarose gels (1.5%), cut from gel and purified by QIAquick Gel Purification Kit (QIAGEN). The resulting sets of the purified restriction fragments from both drFP583 and dmFP592 were combined together and 50 ng of them were put into ligation mix (1X T4 DNA ligation buffer, 400 NEB U of T4 DNA ligase) in total volume of 30 ml. The ligation was performed for 3h at room temperature and stopped by heating at 70°C within 20 min.

The ligation mixture was then diluted by water ten-folds, and 1 ml of the dilution was preformed for PCR reaction (20 cycles, 95°C, 15 sec. 68°C 1min 20 sec) as template. Four oligonucleotides A, B, C, and D (SEQ ID Nos. 58-61, respectively) were used simultaneously as primers for amplifying these fragments in a concentration of 0.1 mM each. After electrophoresis in an agarose gel (1.5%), the target fragment was purified by QIAquick Gel Purification Kit (QIAGEN) and digested with restriction endonucleases BamHI and XhoI (30-50 U each) simultaneously in BamH I restriction buffer (NEB) supplemented with BSA for 3h at 37°C. After purification, the resulting fragment was cloned in pQE-30 plasmid linearized by BamHI and SalI. Ligation reaction was performed in 1X T4 DNA ligation buffer and 400 NEB U of T4 DNA ligase with a total volume of 20 ml for overnight at 16°C. After transformation of *E.coli* cells by 1/5 of the ligation volume and incubation on LB/1% agar plates which were supplemented by 100

mg/ml Ampacilin and 0.1 mM IPTG at 37°C for overnight, the resulting *E.coli* colonies were screened visually under fluorescent microscope using rhodamine filter set. The brightest red colonies were picked up and placed in 200 ml LB medium with 100 mg/ml of Ampacilin. At  
 5 OD<sub>600</sub>=0.6, the *E.coli* culture was induced by IPTG (final concentration was 1 mM) and the fermentation continued for overnight. Purification of recombinant protein containing N-terminus 6Xhis tag was performed using TALON metal-affinity resin according to manufacturer's protocol.

10

### EXAMPLE 15

#### Spectral Properties of drFP583/dmFP592 Hybrid

The emission and excitation spectra for drFP583/dmFP592  
 15 hybrid protein are basically the same as for dmFP592. Table 9 lists the spectral properties of drFP583/dmFP592 hybrid protein.

**TABLE 9**

#### 20 Spectral Properties of drFP583/dmFP592 Hybrid

nFP Name	Absorbance Maximum n m	Emission Maximum n m	Maximum Extinction Coeff.	Relative Quantum Yield*	Relative Brightness **
drFP583/ dmFP592	573	592	35,000	0.24	0.3

\*relative quantum yield was determined as compared to the quantum yield of *A. victoria* GFP.

\*\*relative brightness is extinction coefficient multiplied by quantum  
 25 yield divided by the same value for *A. victoria* GFP.

**EXAMPLE 16**Humanized drFP583/dmFP592 Hybrid and Mutants

drFP583/dmFP592 hybrid was humanized (SEQ ID No. 62 for  
5 nucleotide sequence and SEQ ID No. 63 for deduced amino acid  
sequence). Further, two mutants were generated based on the  
humanized drFP583/dmFP592, i.e., drFP583/dmFP592-2G and  
drFP583/dmFP592-Q3. drFP583/dmFP592-2G contains two  
substitutions, K15Q and T217S (SEQ ID Nos. 64 and 65 for nucleotide  
10 and deduced amino acid sequences, respectively). This mutant was  
derived from the humanized drFP583/dmFP592 hybrid gene by random  
mutagenesis using Diversity PCR Mutagenesis Kit (Clontech) according  
to the corresponding protocol. drFP583/dmFP592-Q3 contains three  
substitutions, K15Q and K83M and T217S (SEQ ID Nos. 66 and 67 for  
15 nucleotide and deduced amino acid sequences, respectively).  
drFP583/dmFP592-Q3 mutant was derived from drFP583/dmFP592-2G  
mutant by random mutagenesis using Diversity PCR Mutagenesis Kit  
(Clontech) according to the corresponding protocol.

drFP583/dmFP592-2G has similar brightness and folding  
20 rate as for non-humanized drFP583/dmFP592 hybrid. While  
drFP583/dmFP592-Q3 could be seen in *E.coli* cells as more dark red  
than parental variant, i.e., drFP583/dmFP592-2G, and the purified  
protein solution has purple color. drFP583/dmFP592-Q3 has the  
emission maximum of 616 nm and excitation maximum of 580 nm.

25

### EXAMPLE 17

#### Possible Applications of Hybrid Mutants

Similar to fluorescent protein drFP583 or dmFP592,  
5 drFP583/dmFP592-Q3 could be used as a tool for investigation of  
protein expression, transport and protein interactions *in vivo*,  
monitoring of promoter activity, and as a transcription reporter or  
fusion tag. Besides, drFP583/dmFP592-Q3 could be chosen as the most  
convenient partner to one of the existing green fluorescent protein  
10 variants in two/triple color labeling assays for simultaneous detection  
of expression of two or more proteins *in vivo* due to its strongly red-  
shifted position of emission maximum and practical absence of  
excitation in green part of spectrum except any spectral overlapping  
and background fluorescence.

15 The method of generating drFP583/dmFP592 hybrid can  
have a general utility for generating hybrid genes (i.e., genes containing  
parts of different genes in various combinations) with improved  
fluorescent characteristics.

Additionally, drFP583/dmFP592-Q3 is the first red-shifted  
20 mutant, which demonstrates that spectral-shifted mutant could be  
obtained by random mutagenesis.

Any patents or publications mentioned in this specification  
are indicative of the levels of those skilled in the art to which the  
invention pertains. These patents and publications are incorporated by  
25 reference to the same extent as if each individual publication was  
specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present  
invention is adapted to carry out the objects and obtain the ends and  
advantages mentioned, as well as those objects and ends inherent

therein. The present examples, along with the methods, procedures, treatments, molecules, and specific compounds described herein, are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes  
5 to the methods and compounds, and other uses, will occur to those skilled in the art and are encompassed within the spirit of the invention as defined by the scope of the claims.



**WHAT IS CLAIMED IS:**

1. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from a Class Anthozoa and wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

10 (c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

15 2. The DNA sequence of claim 1, wherein said organism is from Sub-class Zoantharia.

20 3. The DNA sequence of claim 2, wherein said organism is from Order Corallimorpharia.

4. The DNA sequence of claim 3, wherein said organism is from Family Discosomatidae.

25 5. The DNA sequence of claim 4, wherein said organism is from Genus Discosoma.

6. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein, wherein said DNA has a sequence selected from the group consisting of  
5 SEQ ID Nos. 55, 57, 62, 64 and 66;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic  
10 code, and which encodes a fluorescent protein.

7. The DNA sequence of claim 6, wherein said DNA encodes a fluorescent protein having an amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.  
15

8. The DNA sequence of claim 6, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.  
20

9. A vector capable of expressing the DNA sequence of claim 1 in a recombinant cell, wherein said vector comprising said DNA and regulatory elements necessary for expression of the DNA in the  
25 cell.

10. The vector of claim 9, wherein said DNA encodes a fluorescent protein having the amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67.

5

11. The vector of claim 9, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

10

12. A host cell transfected with the vector of claim 9, wherein said cell is capable of expressing a fluorescent protein.

15

13. The host cell of claim 12, wherein said cell is selected from the group consisting of bacterial cells, mammalian cells, plant cell, yeast and insect cells.

20

14. The host cell of claim 13, wherein said bacterial cell is an *E. coli* cell.

15. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

25

(a) an isolated DNA which encodes a fluorescent protein from an organism from Class Anthozoa, wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

5

16. The isolated and purified fluorescent protein of claim 15, wherein said organism is from Sub-class Zoantharia.

10

17. The isolated and purified fluorescent protein of claim 16, wherein said organism is from Order Corallimorpharia.

15

18. The isolated and purified fluorescent protein of claim 17, wherein said organism is from Family Discosomatidae.

20

19. The isolated and purified fluorescent protein of claim 18, wherein said organism is from Genus Discosoma.

20. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

(a) isolated DNA which encodes a fluorescent protein having an amino acid sequence selected from the group consisting of SEQ ID Nos. 56, 63, 65 and 67;

(b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) isolated DNA differing from said isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

5

21. The isolated and purified fluorescent protein of claim 20, wherein said protein is drFP583.

22. The isolated and purified fluorescent protein of claim  
10 20, wherein said protein is drFP583/dmFP592 hybrid protein.

23. An amino acid sequence which can be used as a basis  
for designing an oligonucleotide probe for identification of a DNA  
encoding a fluorescent protein by means of hybridization, wherein said  
15 sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8,  
11, 12, 14.

24. The amino acid sequence of claim 23, wherein said  
oligonucleotide has a nucleotide sequence selected from the group  
20 consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.




Figure 1




Figure 2




Fig. 3 A



**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**



Fig. 3 B




Fig. 3 c



Fig. 3 D




Fig. 4 F




Fig. 4B




Fig. 4c






Fig. 4 D

## SEQUENCE LISTING

<110> Lukyanov, Sergey A.  
Labas, Yulii A.  
Matz, Mikhail V.  
5 Fradkov, Arcady F.  
Ding, Li  
Chen, Ying  
Green, Gisele

<120> Fluorescent proteins from non-bioluminescent  
10 species of Class Anthozoa, genes encoding such  
proteins and uses thereof

<130> D6196D6/PCT

<140> 09/418,529

<141> 1999-10-14

15 <150> 09/210,330

<151> 1998-12-11

<160> 67

<210> 1

20 <211> 25

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

25 <223> primer TN3 used in cDNA synthesis and RACE

<400> 1

cgcgagtcgac cggtttttttt ttttt 25

<210> 2

30 <211> 23

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

35 <223> primer TS used in cDNA synthesis and RACE

<400> 2

aagcagtggt atcaacgcag agt 23

<210> 3  
 <211> 6  
 <212> PRT  
 <213> *Aequorea victoria*  
 5 <220>  
 <222> 21  
 <223> amino acid sequence of a key stretch on which  
 primer NGH is based; Xaa at position 21  
 represents unknown  
 10 <400> 3  
 Gly Xaa Val Asn Gly His  
 5  
 <210> 4  
 15 <211> 20  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 20 <222> 12  
 <223> primer NGH used for isolation of fluorescent  
 protein; n at position 12 represents any of the  
 four bases  
 <400> 4  
 25 gayggctgcg tnaayggdca 20  
 <210> 5  
 <211> 5  
 <212> PRT  
 30 <213> *Aequorea victoria*  
 <220>  
 <222> 31...35  
 <223> amino acid sequence of a key stretch on which  
 primers GEGa and GEGb are based  
 35 <400> 5  
 Gly Glu Gly Glu Gly  
 5

	<210>	6
	<211>	20
	<212>	DNA
5	<213>	artificial sequence
	<220>	
	<221>	primer_bind
	<223>	primer GEGa used for isolation of fluorescent protein
0	<400>	6

gttacaggtg arggmgargg 20

	<210>	7
	<211>	20
15	<212>	DNA
	<213>	artificial sequence
	<220>	
	<221>	primer_bind
	<223>	primer GEGb used for isolation of fluorescent
20		protein
	<400>	7

gttacaggtg arggkgargg 20

25	<210>	8
	<211>	5
	<212>	PRT
	<213>	<i>Aequorea victoria</i>
	<220>	
30	<222>	31..35
	<223>	amino acid sequence of a key stretch on which primers GNGa and GNGb are based
	<400>	8

Gly Glu Gly Asn Gly  
5

35		
	<210>	9
	<211>	20

<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
5 <223> primer GNGa used for isolation of fluorescent  
protein  
<400> 9

gttacaggtg arggmaaygg 20

10 <210> 10  
<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
15 <221> primer\_bind  
<223> primer GNGb used for isolation of fluorescent  
protein  
<400> 10

gttacaggtg arggkaaygg 20

20  
<210> 11  
<211> 5  
<212> PRT  
<213> *Aequorea victoria*  
25 <220>  
<222> 127..131  
<223> amino acid sequence of a key stretch on which  
primer NFP is based  
<400> 11

30 Gly Met Asn Phe Pro  
5

<210> 12  
<211> 5  
35 <212> PRT  
<213> *Aequorea victoria*  
<220>

<222> 127..131

<223> amino acid sequence of a key stretch on which  
primer NFP is based

<400> 12

5 Gly Val Asn Phe Pro

5

<210> 13

<211> 20

10 <212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

15 <223> primer NFP used for isolation of fluorescent  
protein

<400> 13

ttccayggtr tgaayttycc

20

<210> 14

20 <211> 4

<212> PRT

<213> *Aequorea victoria*

<220>

<222> 134..137

25 <223> amino acid sequence of a key stretch on which  
primers PVMa and PVMb are based

<400> 14

Gly Pro Val Met

30

<210> 15

<211> 21

<212> DNA

<213> artificial sequence

35 <220>

<221> primer\_bind

<222> 15

<223> primer PVMa used for isolation of fluorescent protein; n at position 15 represents any of the four bases

5 <400> 15

cctgccrayg gtcnngtmat g 21

<210> 16

<211> 21

10 <212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<222> 15

15 <223> primer PVMb used for isolation of fluorescent protein; n at position 15 represents any of the four bases

<400> 16

cctgccrayg gtcnngtkat g 21

20

<210> 17

<211> 47

<212> DNA

<213> artificial sequence

25 <220>

<221> primer\_bind

<223> primer T7-TN3 used in cDNA synthesis and RACE

<400> 17

gtaatacgac tcactatagg gccgcagtcg accgtttttt ttttttt 47

30

<210> 18

<211> 45

<212> DNA

<213> artificial sequence

35 <220>

<221> primer\_bind

<223> primer T7-TS used in cDNA synthesis and RACE

&lt;400&gt; 18

gtaatacgac tcactatagg gcaagcagtg gtatcaacgc agagt

45

&lt;210&gt; 19

5 &lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

10 &lt;223&gt; primer T7 used in cDNA synthesis and RACE

&lt;400&gt; 19

gtaatacgac tcactatagg gc

22

&lt;210&gt; 20

15 &lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

20 <223> gene-specific primer used for 5'-RACE for  
*Anemonia majano*

&lt;400&gt; 20

gaaatagtca ggcatactgg t

21

25 &lt;210&gt; 21

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

30 &lt;221&gt; primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Anemonia majano*

&lt;400&gt; 21

gtcaggcata ctggtaggat

20

35



<210> 22  
<211> 21  
<212> DNA  
<213> artificial sequence  
5 <220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Clavularia sp.*  
<400> 22

10 cttgaaatag tctgctatat c 21

<210> 23  
<211> 19  
<212> DNA  
15 <213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Clavularia sp.*  
20 <400> 23

tctgctatat cgtctgggt 19

<210> 24  
<211> 23  
25 <212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
30 *Zoanthus sp.*  
<400> 24

gttcttgaaa tagtctacta tgt 23

<210> 25  
35 <211> 20

<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
5 <223> gene-specific primer used for 5'-RACE for  
*Zoanthus sp.*  
<400> 25  
gtctactatg tcttgaggat 20

10 <210> 26  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
15 <221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma sp.* "red"  
<400> 26  
caagcaaattg gcaaaggatc 19

20  
<210> 27  
<211> 19  
<212> DNA  
<213> artificial sequence  
25 <220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma sp.* "red"  
<400> 27

30 cggattattgtg gccttcgta 19  
  
<210> 28  
<211> 19  
<212> DNA  
35 <213> artificial sequence  
<220>

<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma striata*  
<400> 28

5 ttgtcttctt ctgcacaac 19

<210> 29

<211> 17

<212> DNA

10 <213> artificial sequence

<220>

<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Discosoma striata*

15 <400> 29

ctgcacaacg ggtccat 17

<210> 30

<211> 20

20 <212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Anemonia sulcata*

25

<400> 30

cctctatctt catttcctgc 20

<210> 31

30 <211> 20

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

35 <223> gene-specific primer used for 5'-RACE for  
*Anemonia sulcata*

<400> 31

tatcttcatt tcctgcgtac

20

5           <210>       32  
            <211>       19  
            <212>       DNA  
            <213>       artificial sequence  
            <220>  
            <221>       primer\_bind  
10           <223>       gene-specific primer used for 5'-RACE for  
            *Discosoma sp. "magenta"*  
            <400>       32

ttcagcaccc catcacgag

19

15           <210>       33  
            <211>       19  
            <212>       DNA  
            <213>       artificial sequence  
            <220>  
20           <221>       primer\_bind  
            <223>       gene-specific primer used for 5'-RACE for  
            *Discosoma sp. "magenta"*  
            <400>       33

acgctcagag ctggggtcc

19

25           <210>       34  
            <211>       22  
            <212>       DNA  
            <213>       artificial sequence  
            <220>  
30           <221>       primer\_bind  
            <223>       gene-specific primer used for 5'-RACE for  
            *Discosoma sp. "green"*  
            <400>       34

ccctcagcaa tccatcacgt tc

22

35

&lt;210&gt;       35

<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
5 <221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma sp. "green"*  
<400> 35  
attatctcag tggatgggtc 20  
10  
<210> 36  
<211> 31  
<212> DNA  
<213> artificial sequence  
15 <220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding  
region of nFPs from *Anemonia majano*  
<400> 36  
20 acatggatcc gctctttcaa acaagtttat c 31  
<210> 37  
<211> 34  
<212> DNA  
25 <213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Anemonia majano*  
30 <400> 37  
tagtactcga gcttattcgt atttcagtga aatc 34  
<210> 38  
<211> 29  
35 <212> DNA  
<213> artificial sequence  
<220>

<221> primer\_bind  
<223> upstream primer used to obtain full coding  
region of nFPs from *Clavularia sp.*  
<400> 38

5 acatggatcc aacatttttt tgagaaacg 29

<210> 39  
<211> 28  
<212> DNA  
10 <213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding  
region of nFPs from *Clavularia sp.*  
15 <400> 39

acatggatcc aaagctctaa ccaccatg 28

<210> 40  
<211> 31  
20 <212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
25 region of nFPs from *Clavularia sp.*  
<400> 40

tagtactcga gcaacacaaa ccctcagaca a 31

<210> 41  
30 <211> 28  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
35 <223> upstream primer used to obtain full coding  
region of nFPs from *Zoanthus sp.*

<400> 41

acatggatcc gctcagtcaa agcacggt 28

<210> 42

5 <211> 32

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

10 <223> downstream primer used to obtain full coding  
region of nFPs from *Zoanthus sp.*

<400> 42

tagtactcga gggtggaact acattcttat ca 32

15 <210> 43

<211> 31

<212> DNA

<213> artificial sequence

<220>

20 <221> primer\_bind

<223> upstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "red"

<400> 43

acatggatcc aggtcttcca agaatgttat c 31

25

<210> 44

<211> 29

<212> DNA

<213> artificial sequence

30 <220>

<221> primer\_bind

<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "red"

<400> 44

35 tagtactcga ggagccaagt tcagcctta 29

<210> 45  
<211> 28  
<212> DNA  
<213> artificial sequence  
5 <220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding  
region of nFPs from *Discosoma striata*  
<400> 45  
10 acatgggatcc agttggtcca agagtgtg 28  
  
<210> 46  
<211> 28  
<212> DNA  
15 <213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma striata*  
20 <400> 46  
tagcgagctc tatcatgcct cgtcacct 28  
  
<210> 47  
<211> 31  
25 <212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding  
30 region of nFPs from *Anemonia sulcata*  
<400> 47  
acatgggatcc gcttcctttt taaagaagac t 31  
  
<210> 48  
35 <211> 28  
<212> DNA  
<213> artificial sequence



<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Anemonia sulcata*  
5 <400> 48  
tagtactcga gtccttggga gcggttg 28  
  
<210> 49  
<211> 30  
10 <212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding  
15 <400> 49  
acatggatcc agttgttcca agaatgtgat 30  
  
<210> 50  
20 <211> 26  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
25 <223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "magenta"  
<400> 50  
tagtactcga ggccattacg ctaatc 26  
  
30 <210> 51  
<211> 31  
<212> DNA  
<213> artificial sequence  
<220>  
35 <221> primer\_bind  
<223> upstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "green"

&lt;400&gt; 51

acatggatcc agtgactta aagaagaaat g

31

&lt;210&gt; 52

5 &lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

10 <223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "green"

&lt;400&gt; 52

tagtactcga gattcggttt aatgccttg

29

15 &lt;210&gt; 53

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

20 &lt;221&gt; primer\_bind

&lt;223&gt; TS-oligo used in cDNA synthesis and RACE

&lt;400&gt; 53

aagcagtgggt atcaacgcag agtacgcrgr grg

33

25 &lt;210&gt; 54

&lt;211&gt; 238

&lt;212&gt; PRT

<213> *Aequorea victoria*

&lt;220&gt;

30 &lt;223&gt; amino acid sequence of GFP

&lt;400&gt; 54

Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu

5

10

15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser

35

20

25

30

Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys

		35		40		45
	Phe Ile Cys Thr	Thr Gly Lys Leu Pro	Val Pro Trp Pro Thr	Leu		
		50		55		60
	Val Thr Thr Phe	Ser Tyr Gly Val Gln	Cys Phe Ser Arg Tyr	Pro		
5		65		70		75
	Asp His Met Lys	Gln His Asp Phe Phe	Lys Ser Ala Met Pro	Glu		
		80		85		90
	Gly Tyr Val Gln	Glu Arg Thr Ile Phe	Phe Lys Asp Asp Gly	Asn		
		95		100		105
10	Tyr Lys Thr Arg	Ala Glu Val Lys Phe	Glu Gly Asp Thr Leu	Val		
		110		115		120
	Asn Arg Ile Glu	Leu Lys Gly Ile Asp	Phe Lys Glu Asp Gly	Asn		
		125		130		135
	Ile Leu Gly His	Lys Leu Glu Tyr Asn	Tyr Asn Ser His Asn	Val		
15		140		145		150
	Tyr Ile Met Ala	Asp Lys Gln Lys Asn	Gly Ile Lys Val Asn	Phe		
		155		160		165
	Lys Ile Arg His	Asn Ile Glu Asp Gly	Ser Val Gln Leu Ala	Asp		
		170		175		180
20	His Tyr Gln Gln	Asn Thr Pro Ile Gly	Asp Gly Pro Val Leu	Leu		
		185		190		195
	Pro Asp Asn His	Tyr Leu Ser Thr Gln	Ser Ala Leu Ser Lys	Asp		
		200		205		210
	Pro Asn Glu Lys	Arg Asp His Met Val	Leu Leu Glu Phe Val	Thr		
25		215		220		225
	Ala Ala Gly Ile	Thr His Gly Met Asp	Glu Leu Tyr Lys			
		230		235		

	<210>	55
30	<211>	898
	<212>	DNA
	<213>	<i>Discosoma sp.</i> "red"
	<220>	
	<221>	CDS
35	<223>	cDNA sequence of drFP583
	<400>	55

gtcctcccaa gcagtggat caacgcagag tacgggggag tttcagccag 50  
 tgacggtcag tgacagggtg agccacttgg tataccaaca aaatgaggtc 100

	<210>	56	
	<211>	225	
20	<212>	PRT	
	<213>	<i>Discosoma</i> sp. "red"	
	<220>		
	<223>	amino acid sequence of drFP583	
	<400>	56	
25	Met Arg Ser Ser	Lys Asn Val Ile Lys Glu Phe Met Arg Phe Lys	
		5 10 15	
	Val Arg Met Glu	Gly Thr Val Asn Gly His Glu Phe Glu Ile Glu	
		20 25 30	
	Gly Glu Gly Glu	Gly Arg Pro Tyr Glu Gly His Asn Thr Val Lys	
30		35 40 45	
	Leu Lys Val Thr	Lys Gly Gly Pro Leu Pro Phe Ala Trp Asp Ile	
		50 55 60	
	Leu Ser Pro Gln	Phe Gln Tyr Gly Ser Lys Val Tyr Val Lys His	
		65 70 75	
35	Pro Ala Asp Ile	Pro Asp Tyr Lys Lys Leu Ser Phe Pro Glu Gly	
		80 85 90	
	Phe Lys Trp Glu	Arg Val Met Asn Phe Glu Asp Gly Gly Val Val	

		95		100		105
	Thr Val Thr Gln Asp Ser Ser Leu Gln Asp Gly Cys Phe Ile Tyr					
		110		115		120
	Lys Val Lys Phe Ile Gly Val Asn Phe Pro Ser Asp Gly Pro Val					
5		125		130		135
	Met Gln Lys Lys Thr Met Gly Trp Glu Ala Ser Thr Glu Arg Leu					
		140		145		150
	Tyr Pro Arg Asp Gly Val Leu Lys Gly Glu Ile His Lys Ala Leu					
		155		160		165
10	Lys Leu Lys Asp Gly Gly His Tyr Leu Val Glu Phe Lys Ser Ile					
		170		175		180
	Tyr Met Ala Lys Lys Pro Val Gln Leu Pro Gly Tyr Tyr Tyr Val					
		185		190		195
	Asp Ser Lys Leu Asp Ile Thr Ser His Asn Glu Asp Tyr Thr Ile					
15		200		205		210
	Val Glu Gln Tyr Glu Arg Thr Glu Gly Arg His His Leu Phe Leu					
		215		220		225

	<210>	57
20	<211>	695
	<212>	DNA
	<213>	artificial sequence
	<220>	
	<223>	nucleotide sequence of humanized drFP583
25	<400>	57

	ggggtaccca	tgcgctcctc	caagaacgtc	atcaaggagt	tcatgcgctt	50
	caaggtgcgc	atggagggca	ccgtgaacgg	ccacgagttc	gagatcgagg	100
	gcgagggcga	gggccgcccc	tacgagggcc	acaacaccgt	gaagctgaag	150
	gtgaccaagg	gcggccccct	gcccttcgcc	tgggacatcc	tgtcccccca	200
30	gttccagtac	ggctccaagg	tgtacgtgaa	gcaccccgcc	gacatccccg	250
	actacaagaa	gctgtccttc	cccgaggggt	tcaagtggga	gcgcgtgatg	300
	aacttcgagg	acggcgggcg	ggtgaccgtg	accaggact	cctccctgca	350
	ggacggctgc	ttcatctaca	aggtgaagtt	catcggcggtg	aacttcccct	400
	ccgacggccc	cgtgatgcag	aagaagacca	tgggctggga	ggcctccacc	450
35	gagcgcctgt	acccccgcga	cggcgtgctg	aagggcgaga	tccacaaggc	500
	cctgaagctg	aaggacggcg	gccactacct	ggtggagttc	aagtccatct	550
	acatggccaa	gaagcccgtg	cagctgcccg	gctactacta	cgtggactcc	600

aagctggaca tcacctccca caacgaggac tacaccatcg tggagcagta 650  
 cgagcgcacc gagggccgcc accacctgtt cctgtaatct agagc 695

	8<210>	58	
5	<211>	31	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
10	<223>	oligonucleotide primer used for amplifying non-humanized wild type coding region fragments from drFP583 and dmFP592	
	<400>	58	
	acatggatcc aggtcttcca agaatgttat c		31
15	<210>	59	
	<211>	26	
	<212>	DNA	
	<213>	artificial sequence	
20	<220>		
	<221>	primer_bind	
	<223>	oligonucleotide primer used for amplifying non-humanized wild type coding region fragments from drFP583 and dmFP592	
25	<400>	59	
	tagtactcga gccaaagttca gcctta		26
	<210>	60	
	<211>	30	
30	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
35	<223>	oligonucleotide primer used for amplifying non-humanized wild type coding region fragments from drFP583 and dmFP592	
	<400>	60	

acatggatcc agttgttcca agaatgtgat

30

<210> 61

<211> 27

5 <212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

10 <223> oligonucleotide primer used for amplifying  
non-humanized wild type coding region  
fragments from drFP583 and dmFP592

<400> 61

tagtactcga ggccattacc gctaatac

27

15 <210> 62

<211> 678

<212> DNA

<213> artificial sequence

<220>

20 <223> nucleotide sequence of humanized  
drFP583/dmFP592

<400> 62

atgagctgca gcaagaacgt gatcaaggag ttcattgcggt tcaaggtgcg 50

gatggagggc accgtgaacg gccacgaggt cgagatcaag ggcgagggcg 100

25 agggccggcc ctacgagggc cactgcagcg tgaagctcat ggtgaccaag 150

ggcgggcccc tccccttcgc cttcgacatc ctcagcccc agttccagta 200

cggcagcaag gtgtacgtga agcaccgcc cgacatcccc gactacaaga 250

agctcagctt ccccgagggc ttcaagtggg agcgggtgat gaacttcgag 300

gacggcggcg tgggtgaccgt gagccaggac agcagcctca aggacggctg 350

30 cttcatctac gaggtgaagt tcattcggcg gaacttcccc agcgacggcc 400

ccgtgatgca gcggcggacc cggggctggg aggccagcag cgagcggctc 450

tacccccggg acggcgtgct caagggcgac atccacatgg ccctccggct 500

cgagggcggc ggccactacc tcgtggagtt caagagcatc tacatggcca 550

agaagccgt gcagctcccc ggctactact acgtggacag caagctcgac 600

35 atcaccagcc acaacgagga ctacaccatc gtggagcagt acgagcggac 650

cgagggccgg caccacctct tcctctga 678

<210> 63

<211> 225  
 <212> PRT  
 <213> artificial sequence  
 <220>

5 <223> amino acid sequence of humanized  
 drFP583/dmFP592

<400> 63

	Met	Ser	Cys	Ser	Lys	Asn	Val	Ile	Lys	Glu	Phe	Met	Arg	Phe	Lys	
					5					10						15
10	Val	Arg	Met	Glu	Gly	Thr	Val	Asn	Gly	His	Glu	Phe	Glu	Ile	Lys	
					20					25						30
	Gly	Glu	Gly	Glu	Gly	Arg	Pro	Tyr	Glu	Gly	His	Cys	Ser	Val	Lys	
					35					40						45
	Leu	Met	Val	Thr	Lys	Gly	Gly	Pro	Leu	Pro	Phe	Ala	Phe	Asp	Ile	
15					50					55						60
	Leu	Ser	Pro	Gln	Phe	Gln	Tyr	Gly	Ser	Lys	Val	Tyr	Val	Lys	His	
					65					70						75
	Pro	Ala	Asp	Ile	Pro	Asp	Tyr	Lys	Lys	Leu	Ser	Phe	Pro	Glu	Gly	
					80					85						90
20	Phe	Lys	Trp	Glu	Arg	Val	Met	Asn	Phe	Glu	Asp	Gly	Gly	Val	Val	
					95					100						105
	Thr	Val	Ser	Gln	Asp	Ser	Ser	Leu	Lys	Asp	Gly	Cys	Phe	Ile	Tyr	
					110					115						120
	Glu	Val	Lys	Phe	Ile	Gly	Val	Asn	Phe	Pro	Ser	Asp	Gly	Pro	Val	
25					125					130						135
	Met	Gln	Arg	Arg	Thr	Arg	Gly	Trp	Glu	Ala	Ser	Ser	Glu	Arg	Leu	
					140					145						150
	Tyr	Pro	Arg	Asp	Gly	Val	Leu	Lys	Gly	Asp	Ile	His	Met	Ala	Leu	
					155					160						165
30	Arg	Leu	Glu	Gly	Gly	Gly	His	Tyr	Leu	Val	Glu	Phe	Lys	Ser	Ile	
					170					175						180
	Tyr	Met	Ala	Lys	Lys	Pro	Val	Gln	Leu	Pro	Gly	Tyr	Tyr	Tyr	Val	
					185					190						195
	Asp	Ser	Lys	Leu	Asp	Ile	Thr	Ser	His	Asn	Glu	Asp	Tyr	Thr	Ile	
35					200					205						210
	Val	Glu	Gln	Tyr	Glu	Arg	Thr	Glu	Gly	Arg	His	His	Leu	Phe	Leu	
					215					220						225



<210> 64  
 <211> 678  
 <212> DNA  
 <213> artificial sequence

5 <220>  
 <223> nucleotide sequence of drFP583/dmFP592-2G  
 <400> 64

```

atgagctgca gcaagaacgt gatcaaggag ttcattgcggt tccaggtgcg 50
gatggagggc accgtgaacg gccacgagtt cgagatcaag ggcgagggcg 100
10 agggccggcc ctacgagggc cactgcagcg tgaagctcat ggtgaccaag 150
ggcgggcccc tccccttcgc cttcgacatc ctcagcccc agttccagta 200
cggcagcaag gtgtacgtga agcaccgccg cgacatcccc gactacaaga 250
agctcagctt ccccgagggc ttcaagtggg agcgggtgat gaacttcgag 300
gacggcgggc tgggtgaccgt gagccaggac agcagcctca aggacggctg 350
15 cttcatctac gaggtgaagt tcattggcgt gaacttcccc agcgacggcc 400
ccgtgatgca gcggcgacc cggggctggg aggccagcag cgagcggctc 450
taccgccggg acggcgtgct caaggcgac atccacatgg ccctccggct 500
cgagggcggc ggccactacc tcgtggagtt caagagcatc tacatggcca 550
agaagcccgt gcagctcccc ggctactact acgtggacag caagctcgac 600
20 atcaccagcc acaacgagga ctacaccatc gtggagcagt acgagcggtc 650
cgagggccgg caccacctct tcctctga 678

```

<210> 65  
 25 <211> 225  
 <212> PRT  
 <213> artificial sequence

<220>  
 <223> amino acid sequence of drFP583/dmFP592-2G

30 <400> 65

```

Met Ser Cys Ser Lys Asn Val Ile Lys Glu Phe Met Arg Phe Gln
                    5                      10                      15
Val Arg Met Glu Gly Thr Val Asn Gly His Glu Phe Glu Ile Lys
                    20                      25                      30
35 Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly His Cys Ser Val Lys
                    35                      40                      45
Leu Met Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Ile
                    50                      55                      60

```

Leu Ser Pro Gln Phe Gln Tyr Gly Ser Lys Val Tyr Val Lys His  
 65 70 75  
 Pro Ala Asp Ile Pro Asp Tyr Lys Lys Leu Ser Phe Pro Glu Gly  
 80 85 90  
 5 Phe Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val  
 95 100 105  
 Thr Val Ser Gln Asp Ser Ser Leu Lys Asp Gly Cys Phe Ile Tyr  
 110 115 120  
 Glu Val Lys Phe Ile Gly Val Asn Phe Pro Ser Asp Gly Pro Val  
 10 125 130 135  
 Met Gln Arg Arg Thr Arg Gly Trp Glu Ala Ser Ser Glu Arg Leu  
 140 145 150  
 Tyr Pro Arg Asp Gly Val Leu Lys Gly Asp Ile His Met Ala Leu  
 155 160 165  
 15 Arg Leu Glu Gly Gly Gly His Tyr Leu Val Glu Phe Lys Ser Ile  
 170 175 180  
 Tyr Met Ala Lys Lys Pro Val Gln Leu Pro Gly Tyr Tyr Tyr Val  
 185 190 195  
 Asp Ser Lys Leu Asp Ile Thr Ser His Asn Glu Asp Tyr Thr Ile  
 20 200 205 210  
 Val Glu Gln Tyr Glu Arg Ser Glu Gly Arg His His Leu Phe Leu  
 215 220 225

<210> 66  
 25 <211> 678  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> nucleotide sequence of drFP583/dmFP592-Q3  
 30 <400> 66

atgagctgca gcaagaacgt gatcaaggag ttcattgcggt tccaggtgcg 50  
 gatggagggc accgtgaacg gccacgagtt cgagatcaag ggcgagggcg 100  
 agggccggcc ctacgagggc cactgcagcg tgaagctcat ggtgaccaag 150  
 ggcgggcccc tccccttcgc cttegacatc ctcagcccc agttccagta 200  
 35 cggcagcaag gtgtacgtga agcaccgcc cgacatcccc gactacatga 250  
 agctcagctt ccccgagggc ttcaagtggg agcgggtgat gaacttcgag 300  
 gacggcgggc tggtgaccgt gagccaggac agcagcctca aggacggctg 350

```

cttcatctac gaggtgaagt tcatcggcgt gaacttcccc agcgacggcc 400
ccgtgatgca gcggcggacc cggggctggg aggccagcag cgagcggctc 450
tacccccggg acggcgtgct caagggcgac atccacatgg ccctccgget 500
cgagggcggc ggccactacc tcgtggagtt caagagcatc tacatggcca 550
5 agaagcccggt gcagctcccc ggctactact acgtggacag caagctcgac 600
atcaccagcc acaacgagga ctacaccatc gtggagcagt acgagcggtc 650
cgagggcgcg caccacctct tcctctga 678

```

```

<210> 67
10 <211> 225
    <212> PRT
    <213> artificial sequence
    <220>
    <223> amino acid sequence of drFP583/dmFP592-Q3
15 <400> 67
    Met Ser Cys Ser Lys Asn Val Ile Lys Glu Phe Met Arg Phe Gln
        5 10 15
    Val Arg Met Glu Gly Thr Val Asn Gly His Glu Phe Glu Ile Lys
        20 25 30
20 Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly His Cys Ser Val Lys
        35 40 45
    Leu Met Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Ile
        50 55 60
    Leu Ser Pro Gln Phe Gln Tyr Gly Ser Lys Val Tyr Val Lys His
25 65 70 75
    Pro Ala Asp Ile Pro Asp Tyr Met Lys Leu Ser Phe Pro Glu Gly
        80 85 90
    Phe Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val
        95 100 105
30 Thr Val Ser Gln Asp Ser Ser Leu Lys Asp Gly Cys Phe Ile Tyr
        110 115 120
    Glu Val Lys Phe Ile Gly Val Asn Phe Pro Ser Asp Gly Pro Val
        125 130 135
    Met Gln Arg Arg Thr Arg Gly Trp Glu Ala Ser Ser Glu Arg Leu
35 140 145 150
    Tyr Pro Arg Asp Gly Val Leu Lys Gly Asp Ile His Met Ala Leu
        155 160 165

```

	Arg	Leu	Glu	Gly	Gly	Gly	His	Tyr	Leu	Val	Glu	Phe	Lys	Ser	Ile
					170					175					180
	Tyr	Met	Ala	Lys	Lys	Pro	Val	Gln	Leu	Pro	Gly	Tyr	Tyr	Tyr	Val
					185					190					195
5	Asp	Ser	Lys	Leu	Asp	Ile	Thr	Ser	His	Asn	Glu	Asp	Tyr	Thr	Ile
					200					205					210
	Val	Glu	Gln	Tyr	Glu	Arg	Ser	Glu	Gly	Arg	His	His	Leu	Phe	Leu
					215					220					225

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29473

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : [IPC 7]: C07K 14/435; C12N 1/00, 1/10, 5/10, 15/12, 15/63

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 252.3, 252.33, 324, 410, 254.11, 348, 369, 69.1; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	MATZ et al. Fluorescent proteins from nonbioluminescent Anthozoa species. Nature Biotechnology. October 1999, Volume 17, No. 10, pages 969-973, see entire document.	1-24
X, P	DE 197 18 640 A1 (WIEDENMANN) 22 July 1999 (22.07.99), see entire document.	15, 20
A	US 5,491,084 A (CHALFIE et al) 13 February 1996 (13.02.96).	15, 20
X	ANDERLUH et al. Cloning, sequencing, and expression of equinatoxin II. Biochemical and Biophysical Research Communications. 1996, Volume 220, No. 2, pages 437-442, see entire document.	1,6, 9, 11-15, 20

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 MARCH 2000

Date of mailing of the international search report

**18 APR 2000**

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GABRIELE ELISABETH BUGAISKY

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29473

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — L	MACEK et al. Intrinsic tryptophan fluorescence of equinatoxin II, a pore-forming polypeptide from the sea anemone, <i>Actinia equina</i> L, monitors its interaction with lipid membranes. <i>European Journal of Biochemistry</i> . 1995, Volume 234, pages 329-335, entire document. Cited as "L" document because it establishes fluorescence of equinatoxin II.	15, 20 — 1, 6, 9, 11-14

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29473

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 6-8, 10, 20-24  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Since the sequence diskette (CRF) submitted by applicant is defective, a sequence search could not be performed. Accordingly, claims 6-8, 10 and 20-24 were searched only in-part, based on a word search.
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/29473

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/320.1, 252.3, 252.33, 324, 410, 254.11, 348, 369, 69.1; 530/350; 536/23.5

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Dialog files 155, 5, 434, 34, 357, 28, 44, 35, 77(Medline, Biosis, Scisearch, Derwent Biotech Abs., Oceanic Abs., Aquatic Sci. & Fish Abs., Dissertation Abs. Online, Conference Papers Index); STN-CAS files registry, CAPLUS; WEST files USPT, Derwent WPI  
search terms: fluoresc? bioluminesc? protein? polypeptide?, anthoz?, zoanthar?, corallimorph?, discosom?, rhodact? coral? cnidar?, anemon? alga, algae, invert? coelenter?, drfp583, mrssknvik/sqsp, vngh/sqep, gegeg/sqep, gegng/sqep, gmnfp/sqep, gvnfp/sqep, gpvm/sqep